

Petroleum Geology: Science and Practice in the 21st Century. New Ideas and Paradigms

N. P. Zapivalov*

Trofimuk Institute of Petroleum Geology and Geophysics SB RAS, Koptyuga 3, Novosibirsk 630090, Russia

Abstract: Oil-and-gas-saturated stratum is an integral interrelated system of rocks (minerals) and fluids (oil, gas, and water). Being a «living» fluid-rock system, it is subject to the laws of auto regulation. It is necessary that research and practical efforts be focused on study and control of the oilfield «life».

Keywords: Origin of petroleum, «life» of an oilfield, rehabilitation cycles, residual oil, metasomatism.

EXTENDED ABSTRACT

Hydrocarbons are reported to occur in all strata of the Earth's crust and are supposed to occur in the cosmic space, too. As to their genesis, there are various authoritative positions.

The author's conception does not imply adherence to any of the existing oil origin hypotheses.

It is virtually impossible to create a general theory of oil origin. To look at regional oil-source rocks as a sole indispensable prerequisite of oil formation is a wrong position.

Oil-and-gas-saturated stratum is an integral interrelated system of rocks (minerals) and fluids (oil, gas, and water). Oil deposit is a «living» fluid-rock system therefore subject to the laws of spontaneous autoregulation. ***It is necessary that research and practical efforts be focused on study and control of the oilfield «life».***

The equilibrium-breaking threshold of perturbation for such a fluid-saturated system during the oilfield development can be estimated with the value of formation pressure equal to 5-8MPa.

In the process of developing an oilfield, and especially at its active (forced) stage, it is necessary to use ***rehabilitation cycles*** so as to restore the energy potential and filtration properties of the stratum and to facilitate fresh formation of hydrocarbon masses. In the long run, it will provide for an increased longevity of the object, higher final oil recovery, and beneficial effect for the environment. It will also save labour and financial

costs. At such wells and zones, temporary conservation is needed to apply rehabilitation cycles there, after which their renewed development will become possible (for instance, in the Mexican bay, North Sea, or West Siberia).

It is possible to initiate an accelerated technogenic process of metasomatic dolomitisation in carbonate reservoirs, thereby creating or renewing highly-productive foci at such oilfields.

At present, efforts should be focused on the efficient well-targeted development of the active oilfields in order to produce the ***residual (hard-to-extract) oil*** in a soft, sparing and non-damaging way (Improved Oil Recovery).

Otherwise, huge masses of the already approved residual reserves will remain unextracted in the Earth' depths waiting for future «discoveries» of the already discovered oil.

1. BASIC CONCEPTS

In petroleum geology of the 21st century the ideas of modern non-linear dynamics with its concepts of chaos and self-organization begin to dominate.

Hydrocarbons are reported to occur in all strata of the Earth's crust and are supposed to occur in the cosmic space, too. As to genesis of hydrocarbons, there are various authoritative positions.

Owing to his petroleum longevity, the author has left behind the classical organic hypothesis, although it was his personal discovery in 1958 when a specific oil-source rock («Bazhenov suite») was first distinguished in the Upper Jurassic system of West Siberia, based on the study of test wells. It was stated in the author's Ph.D. thesis in 1962. Nevertheless, now the author

*Address correspondence to this author at Trofimuk Institute of Petroleum Geology and Geophysics SB RAS, Koptyuga, 3, Novosibirsk 630090, Russia; Tel: 7 903 935 87 25; E-mail: ZapivalovNP@ipgg.sbras.ru

holds to a different scientific paradigm, which does not imply adherence to any of the existing oil origin hypotheses. «Road to the Truth passes through a court martial of Thought sitting in session incessantly inside one's mind» (Olzhas Suleimenov, a Kazakh writer and diplomat).

It is virtually impossible to create a general theory of oil origin. To look at regional oil-source rocks as a sole indispensable prerequisite of oil formation is a wrong position.

As early as in the middle of the 20th century, A.I. Levorsen arrived at the conclusion that oil-source rocks have no concern with actual prospecting and exploration works. In his *Geology of Petroleum* [1] he stated: «The problem of oil-and-gas origin now loses its significance as a necessary prerequisite in planning the exploration works. The reason is that petroleum-like hydrocarbons are discovered in almost any non-reservoir rocks. The amount of residual oil [micro-oil] dispersed in these rocks exceeds all the proved reserves in the Earth. Therefore, **there is no need in search for special source rocks**» (in a reverse translation from the Russian edition: Publishing House «Mir», Moscow, 1970, «Conclusion», p. 488).

Various kinds of paleo-reconstructions can hardly be considered a reliable guide in choosing favorable petroleum objects, because each fluid-rock system is subject to secondary, superimposed processes. It is especially important to take into account metasomatic processes. This is why we often deal with young deposits and modern porosity-and-permeability characteristics of a reservoir; their transformation may proceed very rapidly. The age-analogy principle and the exemplar-based method for evaluation of reserves' categories do not meet the reliability requirements in solving problems of petroleum geology. N.A. Eremenko and G. Chilingar [2] took up the position that in a very short geological time reservoir may turn into a caprock and a caprock into a reservoir.

Oil-and-gas accumulations are traced in all types of rocks and at all stratigraphic levels both onshore and offshore. In fact, we live in a hydrocarbon civilization. Our planet is actually a large overall petroleum polygon.

New evidence and facts refuted the myth of «Peak Oil». Oilfields can be discovered in most unexpected places and conditions. The oil recovery rates depend on various natural, technogenic, and market fluc-

tuations including human factor. A great variety of hydrocarbon sources, both conventional and unconventional, have been discovered and are still being discovered; innovative methods and technologies for hydrocarbon production and utilization are being created; on the whole it shows that Peak Oil is dead. **The era of hydrocarbon civilization will never end.**

- Oil-and-gas-saturated stratum is an integral interrelated system of rocks (minerals) and fluids (oil, gas, and water). An oilfield is a «**living**» **fluid-rock system**, subject to the laws of **spontaneous autoregulation**. It is necessary that research and practical efforts be focused on study and **control of the oilfield «life»**. This «life» depends on a great number of rapidly changing factors (gradients); oil deposits may be formed, then dissolve, and after that re-generate again. This is why many of oil-and-gas accumulations are young.
- During the development of an oilfield, the **perturbation threshold** for breaking the equilibrium of this fluid-saturated system can be defined with the value of formation depression. There are strong reasons to estimate the critical value of formation depression as 5-8MPa. This value is almost universal and applicable for all types of reservoirs.
- Fluid-dynamic systems (oilfields, deposits) are in many respects focal and in all likelihood fractal objects with non-uniform productivities.
- As a first approximation, dynamics of the states of a hydrocarbon deposit can be described with an evolutionary equation.

$$\frac{\partial \mathbf{p}}{\partial t} = Z(\mathbf{p}, \mathbf{a}, t),$$

where $\mathbf{p} = (p_1, p_2, \dots, p_k)$ is a chosen set of dynamic variables describing the system state, $\mathbf{a} = (a_1, a_2, \dots, a_l)$ is a set of the system's parameters, t is the time variable, and Z is a non-linear operator of the said variables \mathbf{p} , universal for all types (or at least for a sufficiently wide class) of oilfields, which is to be determined on the basis of empirical regularities of the evolution of hydrocarbon accumulations.

This is the usual form for the dynamic equations in such spheres as dynamics of mechanical systems, fluid dynamics, theory of classical physical fields, atomic systems, evolution of galaxies etc.; in all these fields,

equations of this type make the basis of a successful research of most sophisticated processes and phenomena.

The equations of transfer and balance of hydrocarbon masses and accompanying materials seem to be the most appropriate basis for constructing fluid-dynamic models. It is especially important in proving the empirical law of critical state threshold for oil-and-gas-saturated systems equal to formation depression of 5-8MPa.

2. PRACTICAL RECOMMENDATIONS

At present, forced commercial development of oilfields is being practiced all over the world in order to quickly produce easy-to-extract oil with the use of all available super-intensive methods and technologies. As a result, the **residual oil** (reserves) now amounts to 55-70%. To extract this residual (hard-to-extract) oil from the productive strata, conceptually new ideas and technologies are required.

In petroleum geology, both research and practice (prospecting and developing works) are to be guided by locating zones (foci) of active gradient fluid behavior at each specific moment of time. Nowadays a lot of such innovative techniques are available, including aerial and space survey. The **DFM-method** developed by V.B. Pisetsky [3] and the **SAM-technology** (spectral analysis of microseisms) by G.V. Vedernikov [4] are highly recommended. These technologies make it possible to distinctly determine the highly productive foci during the prospecting and developing works.

In the process of developing an oilfield and especially at its active stage, it is necessary to use **rehabilitation cycles**, in order to restore the energy potential of the stratum and filtration properties of the rock as well as to facilitate fresh formation of hydrocarbon masses. In the long run it will provide for an increased longevity of the object, higher **final oil recovery**, and beneficial effect for the environment.

Due to various reasons, many wells, separate zones and oilfields have to be abandoned. At such wells and zones, temporary conservation should be performed in order to apply rehabilitation cycles. After that, their renewed development will become possible (for instance, in the Mexican bay, North Sea, West Siberia etc.). This idea has been confirmed in a recent article by I.A. Dyachuk [5]. However re-formation of deposits according to the principle of capillary-gravity

segregation seems to be a simplified model: natural processes are much more complicated.

Dolomites in carbonate rock masses formed through late metasomatism are of a special interest. Nano-scale metasomatic processes not only increase porosity and permeability but also often facilitate formation of prolific carbonate reservoirs. It is possible to initiate an accelerated technogenic process of **metasomatic dolomitisation** in order to create or renew highly-productive foci at an oilfield.

It is highly desirable to organize **research-and-development polygons** in various petroleum regions, like the one at the giant Eugene Island oilfield in the Mexican bay (GBRN, Global Basin Research Network).

Regular observations are required with constant monitoring of the current changes inside the fluid-saturated stratum. «Thousands of observations and thousands of measurements are needed to avoid a single error» (Omar Khayyam, a mediaeval astronomer and philosopher). It should be kept in mind that for the Kola Super deep Borehole in Russia (12 262 meters) many preliminary geological and geophysical models proved inadequate.

Virtual models of various kinds (geochemical, seismic, mathematical etc.) are unable to represent adequately the dynamics of oilfield life. George E. P. Box, a renowned specialist in mathematical statistics, remarked: «In fact all models are wrong but some are useful» [6].

The same was clearly shown by Xiao-Hui Wu, Senior Earth Modeling Advisor at ExxonMobil Upstream Research Company, in his work «How to Predict Reservoir Performance with Subsurface Uncertainty at Multiple Scales?». He asserts: «It is possible to reduce the source of uncertainty by reducing modeling and numerical errors with the use of parameters based on field data » [7].

Geofluid-dynamic monitoring of the Earth's depths lags much behind cosmic monitoring, and this retardation may prove fatal for civilization.

3. TODAY'S PROBLEMS

The **era of big oil** in the abyssal Paleozoic of West Siberia [6] is not finished yet. The author has devoted much effort to solving this problem. A. A. Trifimuk, an outstanding Russian petroleum geologist, used to say that the Paleozoic is «a golden substrate of West Siberia».

In the foreseeable future, the task of large-scale industrial development of Russian Arctic off-shore petroleum resources, which are of merely prognostic character, will have to face insurmountable difficulties and can only help in solving some geopolitical problems whereas the vast on-shore transpolar territory of West Siberia is a quite another, already half-solved problem.

Bazhenites and shale oil in West Siberia may look inspiring but in fact no breakthrough can be expected here in near future.

According to recent data [5], sediments of the Bazhenov formation geochemically belong to metal-bearing black shales; as compared with the world-average values for black shales, the Bazhenov formation is three times richer in U, Zn, Sr, Ba and 1.5 times richer in As, Co, and Tb. The concentration of uranium in the rocks is 2 to 171g/ton, with the average value 40.9g/ton. The resources of uranium in this region can be approximately estimated as 3 billion tons.

So, a reasonable question arises whether it may prove commercially viable to produce uranium and other metals from West-Siberian bazhenites, at least in separate promising regions (maybe together with oil).

4. MAIN CONCLUSION

At present, efforts should be focused, first, on the efficient well-targeted development of the active oilfields in order to produce the residual (hard-to-extract) oil in a soft, sparing and non-damaging way (Improved Oil Recovery) and second, on discovering new hydrocarbon accumulations, including secondary ones, throughout the whole stratigraphic section, with all depth horizons and various rock-fluid associations, in regions where a well-developed diversified infrastructure is already available.

Otherwise, huge masses of the already approved residual reserves will remain in the Earth' depths waiting for future «discoveries» of the already discovered oil.

To solve these problems, the whole potential of the powerful Russian science is required; the management of subsurface resources must be science-intensive.

*These ideas, conceptions, paradigms, and suggestions are expressed in many author's publications and public reports of the recent 15 years. Some of them:

MONOGRAPHS

1. Zapivalov NP and Popov IP. Fluid-dynamic models of oil-and-gas deposits. – Novosibirsk: Geo 2003; 198. (*in Russian*)
2. Zapivalov NP, Smirnov GI and Kharitonov VI. Fractals and nanostructures in petroleum geology and geophysics. – Novosibirsk: Geo 2009; 131. (*in Russian*)
3. Zapivalov NP. Petroleum Geology and Geophysics in the 21st Century. A compendium of scientific works published in DEW Journal over thirteen years. – Monograph: Technology publications: Technical books publishers and contributors, 2014. – Editors: Mikhail Eпов, Arun Kumar Singhal. – Dehradun, India–244 p.
4. Zapivalov NP. Offshore oil and gas of the World. –Novosibirsk: Novosibirsk State University 2009; 260. (*in Russian*)

ARTICLES

5. Zapivalov NP. Dynamics of oilfield life // Academy Publish: Journal of Engineering and Technology (*scientific electronic publisher*). ISSN: 2161-7155. 2(1): 112-120. <http://www.academypublish.org/paper/DYNAMIC-S-of-oilfield-life>
6. Zapivalov NP. Rehabilitation cycles as the basis of efficient use of the oil resources // DEW: Drilling and Exploration World. – India 2015; 24(07): 49-53.
7. Zapivalov NP. Rehabilitation cycles as the basis of efficient use of the oil resources // Proc. of the XI International Scientific Congress «Interexpo GEO-Siberia 2015» (April 13-25, 2015). – Novosibirsk: Siberian State University of Geosystems and Technologies 2015; 1: 43-48. (*in Russian*)
8. Zapivalov NP. Metasomatic dolomitization and oil-and-gas content of carbonate rocks // Oil Industry - Neftyanoe Khozyaystvo 2010; 4: 46-49. (*in Russian*)
9. Zapivalov NP. Nano-effects of formation of secondary highly productive reservoirs in carbonate rocks // Georesursy (Georesources), Kazan 2010; 2: 2-6. (*in Russian*)

REFERENCES

- [1] Levorsen AI. Geology of petroleum. Ed. by FAF. Berry WH. Freeman, San Francisco, 2nd ed., 1967; 724 p.
- [2] Eremenko NA and Chilingar GV. Geology of oil and gas at the turn of the century. – Nauka, Moscow 1996; 176 p. (*in Russian*)
- [3] Pisetsky VB. Prognosis of fluid-dynamic parameters of a basin based on seismic data. – Yekaterinburg: Ural State Mining University 2011. (*in Russian*)
- [4] Vedernikov GV. Hydrocarbons pools forecasting by characteristics of microseisms. Novosibirsk: Publishing House «Svin'in i synov'ya», 2012. 201 p. (*in Russian*)
- [5] Dyachuk IA. Reformation of oil fields and reservoirs // Georesursy (Georesources), Kazan, 2015; 1(60): 39-45. (*in Russian*)
- [6] Box GEP and Draper NR. Empirical Model Building and Response Surfaces. John Wiley and Sons, New York, NY 1987; 424.
- [7] Xiao-Hui Wu. How to Predict Reservoir Performance with Subsurface Uncertainty at Multiple Scales? // Lecture. – Society of Petroleum Engineers: Distinguished Lecturer Program 2015.
- [8] Ayzberg RE, Garetsky RG, Zapivalov NP, *et al.* Problems of oil-and-gas capacity of the Upper-Proterozoic and Paleozoic complexes in Belarus and Siberia. Ed. by V.A. Moskvich. BelGeo, Minsk, 2003; 360 p. (*in Russian*)
- [9] Rikhvanov LP, Usoltsev DG, Ilenok SS, *et al.* Mineralogical and geochemical features of the Bazhenov formation, West Siberia, according to nuclear-physics and electron-microscopic methods of research // Bulletin of the Tomsk Polytechnic University, 2015; 326(1): 50-63. (*in Russian*)

Received on 30-09-2015

Accepted on 12-12-2015

Published on 31-12-2015

DOI: <http://dx.doi.org/10.15377/2409-787X.2015.02.02.1>

© 2015 N. P. Zapivalov; Avanti Publishers.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.